

Insect Pest Management in Menthol Mint

Sandeep K^{1*}, Manoj Kumar² and Shashank D U³

¹Department of Entomology, PGCA, Dr. Rajendra Prasad Central Agricultural University, Pusa

²Associate Professor, Department of Entomology, PGCA,

Dr. Rajendra Prasad Central Agricultural University, Pusa

³Ph.D. Scholar, Department of Entomology, PGCA,

Dr. Rajendra Prasad Central Agricultural University, Pusa (Samastipur), Bihar, India

Corresponding Author

Sandeep K

Email: sandeep158k@gmail.com



OPEN ACCESS

Keywords

Insect, Pest management, Integrated Pest Management, Biological control, Botanical insecticides, Sustainable agriculture.

How to cite this article:

Sandeep, K., Kumar, M. and Shashank, D. U. 2024. Insect Pest Management in Menthol Mint. *Vigyan Varta* 5(5): 266-271.

ABSTRACT

The cultivation of Japanese Mint (*Mentha arvensis*) in northern India is a vital practice due to its high menthol content, making it a key source of mint oil for various industries. However, this cultivation faces challenges from insect pests like the mint flea beetle, mint bud mite, mint aphid, cutworms, and thrips, which can significantly impact crop health and yield. To combat these pests sustainably, farmers employ Integrated Pest Management (IPM) strategies, including cultural practices, biological control, chemical control, monitoring, and mechanical methods. By integrating these approaches, farmers can effectively manage pest populations while minimizing environmental impact. Additionally, the use of biological control agents like ladybugs and botanical insecticides derived from mint essential oils offers a sustainable alternative to synthetic pesticides. Implementing these eco-friendly pest management practices is crucial for ensuring the long-term health, yield, and quality of menthol mint crops, contributing to biodiversity conservation and ecosystem balance in mint fields.

INTRODUCTION

Menthol mint cultivation, particularly Japanese Mint (*Mentha arvensis*), is a significant practice with high economic importance. Japanese Mint is extensively cultivated in northern India and is a primary source of menthol, with its essential oil containing around 75-80% menthol content. This mint species is grown as an annual in sub-tropical parts of north India and is propagated through vegetatively. The cultivation of Japanese Mint has been a commercial practice in India for the past four decades, with a substantial production of mint oil. The oil extracted from Japanese Mint is widely used in various industries, including pharmaceuticals, cosmetics, and food flavouring. It is a key ingredient in products like cough drops, mouthwashes, and beverages. The cultivation of menthol mint, especially Japanese Mint, plays a crucial role in the mint industry, meeting the demand for natural menthol preferred in the food and flavour industry (Husain *et al.*, 1988). The cultivation of menthol mint faces a persistent challenge from insect pests, impacting crop productivity. Insect pests pose a significant threat to menthol mint crops, affecting their growth and quality. Various lepidopteran pests infest menthol mint in India, particularly in major growing districts of the Indo-Gangetic plains. These pests cause damage during different stages of crop development, leading to economic losses for mint farmers (Srivastava *et al.*, 2002). The presence of these insect pests necessitates effective pest management strategies to protect menthol mint crops and ensure optimal yields. Research and studies focus on understanding the pest complex infesting menthol mint, highlighting the importance of pest control measures in sustaining the cultivation of this valuable crop (Kedar *et al.*, 2023).

Effective pest management plays a crucial role in maintaining the quality and yield of menthol mint crops. Pest infestations, particularly by lepidopteran pests, can significantly impact the growth and productivity of menthol mint plants, leading to economic losses for farmers. Implementing efficient pest management strategies is essential to mitigate the damage caused by insect pests and ensure optimal crop quality and yield. Research indicates that pests like lepidopteran insects can cause injury to menthol mint plants during various stages of crop development, emphasizing the importance of pest control measures. By effectively managing insect pests, farmers can protect their crops from damage, allowing the plants to grow healthily and produce high-quality mint essential oil. Furthermore, pest management practices, such as scouting for pests, using pesticides when necessary, and implementing integrated pest management techniques, help prevent crop losses and maintain the overall health of menthol mint fields. Controlling pests like the mint flea beetle is crucial to prevent stunted growth, discoloration, and potential plant death, which can impact the yield and quality of the crop (Weller *et al.*, 2000).

Insect Pests of Menthol Mint Crops

The common insect pests affecting menthol mint crops include Mint flea beetle (*Longitarsus waterhousei*) This beetle feeds on the lower leaves in the inner canopy and causes little financial crop damage, but its presence can stunt plant growth and affect the overall health of the mint plants. Mint bud mite (*Floridotarsonemus sp.*) This mite causes stunting and distortion of the upper plant late in the season, with symptoms such as shortened terminal internodes, curling of new leaves, and a twisting or puckering of apical buds, commonly referred to as "squirrelly mint" Spearmint is less severely damaged but

can also sustain high mint bud mite populations. Mint aphid (*Ovatus crataegarius*) Mint aphids are yellow green to apple green, mottled with darker green, and are 1.5 to 2 mm long. They overwinter around the base of mint plants and give birth to living nymphs in the spring. Aphids feed on stems and the undersides of mint leaves, causing discoloration, stunted growth, and curled leaves. Cutworms (*Agrotis spp.*) These can cause damage by severing the stems of young transplants or seedlings at the soil line, or by eating irregular holes into the surface of fruits. Larvae are usually active at night and hide during the day in the soil at the base of the plants. Thrips (*Frankliniella occidentalis*) this can transmit viruses such as Tomato spotted wilt virus and cause damage by feeding on the undersides of leaves, resulting in distorted leaves and reduced oil yields. These pests can significantly impact the growth and productivity of menthol mint crops, leading to economic losses for farmers (Berry *et al.*, 1977)

Insect pests have a significant impact on mint crops, affecting crop health, yield, and quality in various ways. The presence of insect pests like the mint flea beetle and mint bud mite can lead to stunted growth, discoloration, distortion of plant parts, and even plant death, ultimately reducing the overall health of the mint crop. These pests can cause damage to the underground parts of the mint plant, leading to decay and secondary infections from soil microorganisms, which further compromise crop health and yield. The damage caused by insect pests can result in reduced oil yields, with the mint bud mite capable of reducing oil yields by as much as 80% due to its stunting and distortion effects on the upper plant. Additionally, insect pests like the mint aphid can discolor foliage, stunt plant growth, cause curled leaves, and damage buds, impacting the overall quality of the mint crop and potentially affecting the flavor and

aroma of the essential oil produced. In terms of yield, insect pests can weaken a mint stand significantly, leading to economic losses for farmers. If pest infestations build up to excessive levels, they can weaken the crop to the point of complete economic loss, especially if the field has to be removed from production prematurely in the rotation cycle. The damage inflicted by insect pests can reduce the marketable yield of mint crops, affecting the profitability of mint production (Murray *et al.*, 2020).

Pest Management in Menthol Mint

Sustainable and eco-friendly insect pest management practices are crucial for maintaining the health, yield, and quality of menthol mint crops. These practices aim to minimize the use of synthetic pesticides and promote the use of natural methods to control insect pests. Some of the key sustainable insect pest management practices in menthol mint farming include:

- **Companion planting:** Planting mint with other plants that repel insect pests, such as catnip, spearmint, peppermint, and other mint plants, can help keep pests at bay.
- **Beneficial insects:** Encouraging beneficial insects like ladybugs, spiders, wasps, dragonflies, lacewings, and praying mantises to inhabit the mint fields can help control pest populations naturally.
- **Neem oil:** Using neem oil as a natural insecticide and repellent can affect the feeding and reproduction cycles of many garden bugs, including aphids, mealybugs, whiteflies, spider mites, thrips, and caterpillars.
- **Organic sprays:** Making homemade sprays using ingredients like insecticidal soap, garlic, and vinegar can help keep insect pests away from the mint plants.

- **Physical barriers:** Using row covers and other physical barriers can protect mint crops from flying insects.
- **Integrated Pest Management (IPM):** Implementing an IPM approach that includes cultural, biological, chemical, and genetic techniques can help manage pests in a sustainable and environmentally friendly manner.
- **Biopesticides:** Developing and using biopesticides, such as biological nematicides, fungicides, and miticides, can help control pests without relying on synthetic chemicals.
- **Education and training:** Providing training for farmers and agricultural professionals on sustainable pest management practices can help promote the adoption of eco-friendly methods in menthol mint farming.

The implementation of Integrated Pest Management (IPM) strategies in menthol mint farming involves a comprehensive approach to pest control that integrates various techniques to manage insect pests sustainably and effectively. In menthol mint farming, IPM strategies typically include a combination of cultural, biological, chemical, and genetic methods to control pest populations while minimizing environmental impact and promoting long-term sustainability. Key components of IPM strategies in menthol mint farming include:

- **Cultural Practices:** Implementing cultural practices like crop rotation, sanitation, and planting disease-free rootstock to prevent the introduction and spread of pests in mint fields.
- **Biological Control:** Encouraging the presence of beneficial insects, such as ladybugs and parasitic wasps, that

naturally prey on pest populations to help control insect pests in a sustainable manner.

- **Chemical Control:** Using pesticides judiciously and selectively when necessary, following guidelines for application timing and dosage to minimize environmental impact and non-target effects.
- **Monitoring and Scouting:** Regularly monitoring mint fields for pest presence and damage through scouting to assess the need for pest control measures and to determine the effectiveness of implemented strategies.
- **Mechanical Control:** Employing physical barriers, traps, and other mechanical methods to manage pest populations, such as using row covers to protect mint plants from flying insects.

By integrating these diverse approaches, menthol mint farmers can effectively manage insect pests, reduce reliance on synthetic pesticides, protect crop health and yield, and contribute to the overall sustainability of menthol mint farming operations (Pushpangadan and Tewari, 2006).

The utilization of biological control agents and botanical insecticides in menthol mint farming is a sustainable approach to pest management that minimizes the use of synthetic chemicals and promotes environmentally friendly practices. Biological control agents, such as beneficial insects like ladybugs and parasitic wasps, play a crucial role in controlling pest populations naturally in menthol mint fields. These beneficial insects prey on insect pests, helping to keep their populations in check without the need for chemical interventions. Botanical insecticides derived from plant extracts, such as mint essential oils, are also utilized in menthol mint farming for pest

control. Mint essential oils, like those from *Mentha piperita* and *Mentha arvensis*, contain active constituents that exhibit insecticidal properties, making them effective natural pesticides for managing insect pests in agriculture. These botanical insecticides offer a sustainable alternative to synthetic pesticides, reducing the environmental impact of pest control practices in menthol mint cultivation (Nazir et al., 2019).

CONCLUSION

Effective insect pest management is crucial in menthol mint cultivation to ensure the health, yield, and quality of mint crops. By implementing successful pest management interventions like Integrated Pest Management (IPM) practices, utilizing biological control agents, and incorporating botanical insecticides, mint farmers can control pest populations sustainably and minimize the use of synthetic chemicals. These strategies help protect mint crops from damaging pests like the mint flea beetle, mint bud mite, and other insect pests, ensuring optimal growth and productivity of menthol mint plants. Additionally, the use of mint volatiles and companion plants can enhance pest management by attracting beneficial insects and predators, contributing to a balanced ecosystem in mint fields. Adopting sustainable and eco-friendly pest control strategies in menthol mint cultivation is of paramount importance for the long-term health and productivity of mint crops. By implementing practices such as Integrated Pest Management (IPM), utilizing biological control agents, and incorporating botanical insecticides derived from mint essential oils, mint farmers can effectively manage pest populations while minimizing environmental impact and reducing reliance on synthetic chemicals. These sustainable pest control strategies not only help protect mint crops from damaging pests like the mint flea beetle and mint bud

mite but also contribute to biodiversity conservation, soil health, and overall ecosystem balance in mint fields. Additionally, practices like companion planting, physical barriers, and organic manure application promote a holistic approach to pest management that supports the sustainability and resilience of menthol mint cultivation.

REFERENCES

- Berry, R. E., & Fisher, G. C. (1977). Insects on mint.
- Husain, A., Virmani, O. P., Singh, D.V., Singh, A. and Singh, K. (1988). Mint farming in India. CIMAP Farm Bulletin No. 11, 5-6.
- Kedar, S. C., Gupta, A., Shashank, P. R., Navik, O., & Patil, J. (2023). The lepidopteran pest complex infesting menthol mint in India: Distribution during the crop development, species composition and associated parasitoids. *Crop Protection*, 173, 106382.
- Murray, K., Walenta, D. L., Jepson, P. C., & Sandlin, I. (2020). Integrated Pest Management Strategic Plan for Oregon, Washington and Idaho Mint Crops: Summary of a Workshop Held on March 8, 2020, Portland, Oregon. Oregon State University Extension Service.
- Srivastava, R. K., Singh, A. K., Kalra, A., Tomar, V. K. S., Bansal, R. P., Patra, D. D., and Kumar, S. (2002). Characteristics of menthol mint *Mentha arvensis* cultivated on industrial scale in the Indo-Gangetic plains. *Industrial crops and products*, 15(3), 189-198.

Nazir, T., Khan, S., & Qiu, D. (2019). Biological control of insect pest. *Pests Control and Acarology*, 21.

Pushpangadan, P., & Tewari, S. K. (2006). Peppermint. In *Handbook of Herbs and Spices* (pp. 460-481). Woodhead Publishing.

Weller, S., Green Jr, R., Janssen, C., & Whitford, F. (2000). Mint production and pest management in Indiana. PPP-103. West Lafayette, IN: Purdue University Cooperative Extension Service.