# The Cashew Advantage: Unlocking High Incomes on Medinipur's Lateritic Soils

**Debjit Chakraborty* and Tandra Das**

*Division of Agronomy, School of Agriculture and Rural Development, Ramakrishna Mission Vivekananda Educational & Research Institute, Narendrapur Campus, Kolkata, West Bengal – 700103, India*

**Corresponding Author**
Debjit Chakraborty
Email: debjitchakraborty9832@gmail.com

**OPEN ACCESS**

**Keywords**

Cashew (*Anacardium occidentale* L.), Medinipur, red and lateritic soils, high-density planting, farm income

*How to cite this article:*

Chakraborty, D. and Das, T. 2026. The Cashew Advantage: Unlocking High Incomes on Medinipur's Lateritic Soils. *Vigyan Varta* 7 (11): 217-221.

**ABSTRACT**

The sweeping, red landscapes of West Bengal's lateritic zone spanning across the rugged terrains of Medinipur are visually striking, but they have historically been a tough battleground for farmers. With erratic moisture, undulating slopes, and naturally low soil fertility, traditional crops often struggle to deliver a profitable return. But there is a high-value, climate-resilient alternative perfectly designed for this exact environment: the cashew. For years, cashew in these regions was treated as a marginal plantation tree planted and forgotten. Today, that narrative is changing. Cashew is no longer just a survival crop for tough soils; it is a gateway to substantial rural wealth. The red soils of Medinipur naturally provide the well-drained, slightly acidic conditions that cashew trees crave. When this natural geographical advantage is paired with an improved package of agricultural practices, the economic results are extraordinary. The secret to unlocking this "red earth goldmine" lies in moving away from outdated methods. By adopting modern scientific cultivation such as selecting high-yielding grafted varieties suited for Medinipur, implementing high-density planting, strategic canopy pruning, and precision nutrient management, farmers can drastically multiply their nut yield and kernel quality. This article explores how upgrading to smart, scientifically backed farming practices can transform cashew from a neglected backyard tree into a highly profitable commercial enterprise, offering a sustainable and

lucrative future for the farmers of Bengal's lateritic belt.

## INTRODUCTION:

The red and lateritic belt of southwestern West Bengal constitutes an important but challenging agricultural environment. Large parts of the region are characterized by undulating topography, highly weathered soils, low organic matter and deficiencies of major plant nutrients. At the Regional Research Station of Bidhan Chandra Krishi Viswavidyalaya (BCKV), Jhargram, representative upland soils of the red and lateritic zone have been described as coarse textured and acidic, with soil pH generally ranging from 5.0 to 5.5, low organic matter and low available nitrogen, phosphorus and potassium. The uplands are also susceptible to erosion and may have shallow rooting conditions with gravel, stones and morum within the plough layer (Bidhan Chandra Krishi Viswavidyalaya [BCKV], n.d.-a).

These characteristics impose substantial limitations on conventional rain-fed agriculture. Low water-holding capacity, seasonal moisture stress and poor nutrient status can reduce crop productivity, particularly when crop selection and management are not matched with the inherent properties of the soil. However, agricultural suitability is crop-specific. A soil that is restrictive for a nutrient-demanding annual crop may still provide a suitable environment for a perennial crop adapted to relatively acidic, well-drained conditions.

Cashew (*Anacardium occidentale* L.) is particularly relevant in this context. The crop is adapted to a range of soil types, including lateritic soils, provided that drainage is adequate. It does not tolerate prolonged waterlogging and performs well under warm climatic conditions (ICAR–Central Coastal Agricultural Research Institute [ICAR-CCARI], n.d.). The compatibility between cashew requirements and several characteristics of the red and lateritic environment provides a scientific basis for considering cashew as a land-use diversification option in Medinipur. Importantly, this opportunity is supported by regional research rather than by ecological assumptions alone. BCKV has maintained an All India Coordinated Research Project on Cashew at its Regional Research Station, Jhargram, since the early 1980s. Research has included cultivar evaluation, propagation, spacing, nutrient management, canopy management, soil and water conservation, intercropping and high-density planting (BCKV, n.d.-b).

The objective of this article is therefore to examine the potential of cashew cultivation in the lateritic zone of Medinipur by integrating three major dimensions: ecological suitability, improved production technology and economic potential. Particular emphasis is placed on the selection of grafted planting material, high-density orchard systems, strategic canopy management, precision nutrient management and soil-water conservation.

### 1. The Medinipur Advantage: Why Lateritic Soil Can Favour Cashew

### 1.1 From Soil Constraint to Crop Opportunity

The agricultural value of lateritic soil should not be judged solely by its performance under conventional field crops. Lateritic soils are products of intense weathering and leaching and may have low fertility, acidity and limited water-holding capacity. These properties can

constrain many crops, but they do not automatically make the land unsuitable for perennial horticulture.

Cashew has several characteristics that make it compatible with such environments. It can grow on a relatively wide range of soils and is particularly suited to well-drained conditions. Heavy soils with poor drainage are considered unsuitable because cashew is sensitive to waterlogging. This creates an important ecological advantage for well-drained portions of the Medinipur lateritic landscape. Rapid drainage can reduce prolonged saturation of the root zone during the monsoon. However, drainage alone is not sufficient. Excessively rapid drainage combined with shallow soil can increase dry-season moisture stress. Therefore, the ideal site is not simply a "dry" soil but a well-drained soil with adequate effective rooting depth and sufficient moisture availability during critical stages (ICAR-CCARI, n.d.).

### 1.2 Soil Acidity as a Manageable Constraint

The acidic reaction of lateritic soils is another major characteristic of the region. BCKV observations from the Jhargram red and laterite zone indicate soil pH values of approximately 5.0–5.5 in representative upland research areas (BCKV, n.d.-a). Cashew is relatively tolerant of acidic soil environments compared with many crops. Nevertheless, acidity should not be regarded as irrelevant. Strong acidity can affect nutrient availability and may increase the risk of aluminium and manganese toxicity. Phosphorus fixation can also reduce the availability of applied phosphorus. Consequently, the appropriate approach is management rather than neglect. Soil testing should be used to determine the severity of acidity and nutrient limitations. Where necessary, suitable soil amendments, organic matter and balanced fertilization should be incorporated into the orchard nutrient-management programme.

### 1.3 Drainage and Root-Zone Conditions

Cashew requires a well-aerated root environment. According to ICAR-CCARI, soils ranging from sandy to lateritic can be suitable for cashew, whereas heavy clay soils with poor drainage are unsuitable because of waterlogging (ICAR-CCARI, n.d.).

The red and lateritic soils of Medinipur can therefore provide a favourable physical environment where adequate depth and drainage occur. Nevertheless, considerable spatial variability exists within the lateritic landscape. Shallow, gravelly or highly eroded uplands should not be treated in the same way as deeper, better-developed soils. Hence, site-specific land evaluation should precede commercial orchard establishment.

## 2. Regional Evidence: BCKV's Jhargram Research

Long-term research at BCKV's Regional Research Station in Jhargram provides location-specific proof that scientific cultivation works in Paschim Medinipur. Moving beyond traditional low-yield methods, modern trials have identified highly productive cultivars for this specific zone (BCKV, n.d.-b; Poduval, 2015):

- Top Performers: Early studies highlighted H-255, H-303, and M-15/4 for their superior canopy development and nut yield (Poduval, 2015).
- High Yielders: BCKV recently released Bidhan Jhargram-2, which yields over 11 kg of nuts per mature tree (approx. 3.5–4.0 tonnes/hectare) (BCKV, n.d.-c).
- High-Density Ready: Bidhan Bonsai Kaju was identified as ideal for modern high-density planting, yielding 2–2.5 tonnes/hectare (BCKV, n.d.-c).

### 3. The Money Math: Economic Potential

Cashew transforms marginalized, rain-fed land into a high-value asset. A mature, well-managed orchard yielding 400–600 kg of raw cashew nuts per acre can generate ₹40,000–₹90,000 in annual gross revenue (assuming a farm-gate price of ₹100–₹150/kg).

While these numbers are attractive, true profitability requires treating cashew as a long-term commercial investment. Net farm income will depend on upfront establishment time, labor, fertilizers, and pest management. Ultimately, cashew offers a significantly higher and more stable economic return than conventional rain-fed crops on Medinipur's lateritic soils provided farmers use high-yielding grafted varieties and modern management techniques.

### 4. Modernizing Cashew Production: The Scientific Package of Practices

Transitioning from traditional, low-yield cashew plantations to highly profitable commercial orchards requires a systemic shift in agronomy. The foundation of this transformation relies on three critical interventions. Genetics and Orchard Architecture Seedling propagation produces unpredictable yields. Commercial viability demands healthy, grafted plants of regionally proven cultivars. Cultivars like Jhargram-1, Bidhan Jhargram-2, and Bidhan Bonsai Kaju (developed by BCKV) offer genetic uniformity and predictable nut quality (BCKV, n.d.-c).

Furthermore, maximizing land-use efficiency requires high-density planting—but this is not simply planting trees closer together. It is an integrated architectural system. Varieties like Bidhan Bonsai Kaju thrive in high-density setups only when paired with strategic canopy pruning. Unmanaged canopies block light and air, whereas systematic pruning maintains an open, productive structure that optimizes photosynthesis and simplifies harvesting.

Precision Nutrition and Moisture Management Medinipur's lateritic uplands are inherently deficient in organic matter and primary nutrients. Blanket fertilization is ineffective, especially because highly weathered acidic soils suffer from severe phosphorus fixation.

Nutrient application must combine organic amendments (to improve soil structure and microbial activity) with precise mineral fertilization. Based on BCKV's regional trials, an annual application of 500 g N + 250 g $P_2O_5$ + 250 g $K_2O$ per plant is highly effective, though actual doses should be fine-tuned via soil testing. Because these undulating landscapes are prone to erosion and seasonal dry spells, nutrient management must be coupled with rigorous soil and water conservation techniques, such as contour trenching, mulching, and rainwater harvesting in tree basins (Mangalassery *et al*., 2022).

### 5. Constraints and Strategic Expansion

While cashew is climate-resilient and suitable for crop diversification, it should not be planted indiscriminately across all degraded lands. Severe limitations such as extremely shallow soil, exposed bedrock, and lack of supplemental irrigation during establishment will guarantee crop failure. Future expansion requires GIS-based land suitability mapping to match orchards to areas with adequate rooting depth.

Finally, the economic potential of cashew in Medinipur depends on the broader value chain. Increased production must be matched with local processing, grading, and marketing infrastructure. Without local value addition, the true wealth generated by these orchards will leave the region rather than enriching the farmers of the lateritic belt.

### 6. Future Perspective: From Red Soil to Red-Earth Value Chain

Future Perspective: The "Red-Earth" Value Chain

The future of cashew cultivation in Medinipur relies on abandoning the outdated "plant and forget" mindset and embracing knowledge-intensive commercial horticulture. The greatest economic leap will not come from a single magic technology, but from mastering the entire production lifecycle as an integrated system.

To unlock true wealth, farmers and policymakers must connect every link in the chain: meticulous site selection, high-quality grafted saplings, optimized planting density, precision canopy and nutrient management, and rigorous soil-water conservation (Mangalassery *et al*., 2022). But the chain does not stop at the farm gate. Quality harvesting must feed directly into local processing and value-addition. Establishing local processing infrastructure ensures that the premium price of finished, graded cashews stays within the local economy, rather than being exported out as lower-value raw nuts.

## CONCLUSION

For decades, the undulating, moisture-starved lateritic belts of Medinipur have been viewed as agricultural liabilities. Cashew flips this narrative. It thrives precisely where conventional rain-fed crops fail, proving that Medinipur's red soil is not barren it simply needed the right biological match. Turning this ecological compatibility into sustained rural wealth requires a fundamental shift in farming culture. The regional breakthroughs at BCKV's Jhargram research station prove that transitioning to modern, scientifically backed practices is not just theoretical; it works in Medinipur's exact soil conditions (BCKV, n.d.-b; Poduval, 2015). High-yielding grafted cultivars, strategic canopy pruning, and soil-test-based nutrient management must form the core of this transformation. This agricultural shift is not a blind gold rush. Future expansion must remain strategic and evidence-based, reserving orchards for sites with adequate rooting depth rather than indiscriminately planting on exposed bedrock. But for the vast stretches of suitable lateritic uplands, the blueprint is clear. By pairing Medinipur's natural geographical advantage with modern horticultural science and local processing infrastructure, cashew can transform degraded red soils into generational assets driving farm income, creating rural jobs, and redefining the sustainable agricultural future of West Bengal.

## REFERENCES

Bidhan Chandra Krishi Viswavidyalaya. (n.d.-a). AICRP on agroforestry: Regional Research Station, Jhargram—soil and agro-ecological characteristics of the red and laterite zone of West Bengal. BCKV.

Bidhan Chandra Krishi Viswavidyalaya. (n.d.-b). AICRP on cashew: Regional Research Station, Jhargram, Paschim Medinipur, West Bengal. BCKV.

Bidhan Chandra Krishi Viswavidyalaya. (n.d.-c). Crop varieties: Bidhan Jhargram-2 and Bidhan Bonsai Kaju. BCKV.

ICAR–Central Coastal Agricultural Research Institute. (n.d.). Cashew: Soil and climate requirements. Indian Council of Agricultural Research.

Mangalassery, S., Adiga, J. D., Veena, G. L., & Manjesh, G. N. (2022). Soil and water conservation, and irrigation management in cashew. ICAR–Directorate of Cashew Research.

Poduval, M. (2015). Performance of cashew (Anacardium occidentale) cultivars under red and laterite zone of West Bengal. Acta Horticulturae, 1080, 105–110. https://doi.org/10.17660/ActaHortic.2015.1080.18.